




HIV AMONG MEN IN MARYLAND, 2014

AUGUST 2016

New HIV Diagnoses

- Men made up **70.8%** (871) of the **1,230** reported adult/adolescent HIV diagnoses (age 13+) in Maryland in **2014**.
- The proportion of HIV diagnoses in Maryland has **decreased** in men from **85.2%** (in 1985) to **70.8%** (in 2014).
- 70.7%** of the **871** reported HIV diagnoses among men (age 13+) in 2014 were among **non-Hispanic (NH) blacks**, **14.1%** NH white, **9.8%** Hispanic, and **5.4%** other races.
- Most HIV diagnoses in men in 2014 were among **Men who have Sex with Men (MSM)** (**74.3%**).



Characteristics of Men Living with HIV

Exposure Category


- 52.1% MSM**
- 24.0% IDU
- 18.5% HET
- 5.0% MSM/IDU




Race/Ethnicity

- 71.6% NH Black**
- 17.3% NH White
- 6.1% Hispanic
- 5.1% NH Other

- Of the **30,463** total living HIV cases (age 13+) reported in Maryland in 2014, **65.1%** (**19,817**) of those were among **males**. Adult/adolescent living case rates (per 100,000) were **twice** as high in **men** (**825.3**) as compared to **women** (**408.8**). In other words, **1 in 121 men** was reported to be living with HIV in Maryland.



- Total male living HIV case rates (all ages) were **4-8 times higher** among **NH blacks** (**1,763.3 cases per 100,000**) as compared to other races. In other words, **1 in every 56 NH black males** in Maryland was reported to be living with HIV.
- Of the **19,817** total living HIV cases among males in Maryland, about half (**50.1%**) had a **viral load test** result reported in 2014. Among those **9,926** males with a reported test result, **75.7%** had a **suppressed viral load**.



MSM: Men who have Sex with Men | **HET:** Heterosexual Exposure | **IDU:** Injection Drug User